

A counterexample to a conjecture of Akemann and Anderson

Nik Weaver*

Akemann and Anderson made a conjecture about “paving” projections in finite dimensional matrix algebras which, if true, would settle the well-known Kadison-Singer problem. We falsify their conjecture by an explicit sequence of counterexamples.

Let H be a separable Hilbert space, let $B(H)$ be the C^* -algebra of bounded linear operators on H , and let $\mathcal{D}$ be a maximal abelian self-adjoint subalgebra (MASA) of $B(H)$. Does every pure state on $\mathcal{D}$ extend uniquely to a pure state on $B(H)$? In [6] Kadison and Singer answered this question negatively for nonatomic MASAs and speculated that the answer would also be negative in the atomic case. This has come to be known as the Kadison-Singer problem.

A simple application of the Hahn-Banach theorem shows that every pure state on $\mathcal{D}$ has at least one extension to a state on $B(H)$, and it is easy to see that the set of all such extensions is convex and weak* compact. Hence there exist extreme points in this set, i.e., pure state extensions. Thus, pure state extensions always exist, and the issue is uniqueness.

Concretely, we can take H to be $l^2(\mathbf{N})$ and $\mathcal{D}$ to be $l^\infty(\mathbf{N})$, regarded as a subalgebra of $B(H)$ via action by multiplication operators, i.e., diagonal matrices. Since $\mathcal{D}$ is abelian, the pure states on $\mathcal{D}$ are precisely the homomorphisms from $\mathcal{D}$ into $\mathbf{C}$, which are well-known to correspond to ultrafilters on $\mathbf{N}$. (The ultrafilter $\mathcal{U}$ on $\mathbf{N}$ gives rise to the complex homomorphism $\rho_{\mathcal{U}}$ defined by $\rho_{\mathcal{U}}(f) = \lim_{\mathcal{U}} f$.)

Is it not too hard to show that if $\mathcal{U}$ is a fixed (trivial) ultrafilter then $\rho_{\mathcal{U}}$ has a unique pure state extension. G. Reid [7] showed that there exist free (nontrivial) ultrafilters $\mathcal{U}$ that also extend uniquely.

Let $\pi : B(l^2(\mathbf{N})) \rightarrow l^\infty(\mathbf{N})$ be the standard conditional expectation, $\pi(T)(n) = \langle Te_n, e_n \rangle$, where (e_n) is the standard orthonormal basis of $l^2(\mathbf{N})$. J. Anderson [3] showed that if ρ is any pure state on $l^\infty(\mathbf{N})$ then $\pi \circ \rho$ is a pure state on $B(l^2(\mathbf{N}))$. Thus, the question of Kadison and Singer becomes “Are there ever any other (pure) state extensions of ρ ?”

The problem can also be rephrased in terms of “paving” finite matrices whose diagonals are zero. The best possible proof of this reduction is given in [8]. See also [5] for partial positive results and further references.

* Supported by NSF grant DMS-0070634

1. The Akemann-Anderson conjectures. Idea of the counterexample

A different, but related, finite dimensional adaptation of the Kadison-Singer problem was given by Akemann and Anderson in Chapter 7 of their memoir [1]. They used the following notation. $H = \mathbf{C}^n$ is a finite dimensional complex Hilbert space and $\mathcal{D}$ is the algebra of diagonal matrices. For any projection $p \in B(H) = M_n(\mathbf{C})$, let

$$\delta_p = \max\{\langle pe_i, e_i \rangle : 1 \leq i \leq n\}$$

where $\{e_i\}$ is the natural basis of $\mathbf{C}^n$. Recall that a “symmetry” is a self-adjoint unitary.

Akemann and Anderson state several versions of their conjecture. The primary version is the following:

Conjecture A [1, Conjecture 7.1.1]. For any projection $p \in M_n(\mathbf{C})$ there exists a symmetry $s \in \mathcal{D}$ such that $\|psp\| \leq 2\delta_p$.

This is equivalent to asking whether for any projection $p \in M_n(\mathbf{C})$ there is a projection $q \in \mathcal{D}$ such that

$$\max(\|qpq\|, \|(1-q)p(1-q)\|) \leq \frac{1}{2} + \delta_p$$

[1, Proposition 7.4]. This puts the conjecture in the form of a paving question.

If true, Conjecture A would imply a positive solution to the Kadison-Singer problem (i.e., unique extension of pure states on atomic MASAs). The grounds for believing the conjecture include an analogy with known combinatorial results [2, 4] as well as computer experimentation which suggests that counterexamples are rare [1, p. 80]. Nonetheless, we will show that this conjecture is false. A weaker version that remains open, and would still resolve the Kadison-Singer problem, is the following:

Conjecture B [1, Conjecture 7.1.3]. There exist $\gamma, \epsilon > 0$ (independent of n) such that for any projection $p \in M_n(\mathbf{C})$ with $\delta_p < \gamma$ there exists a symmetry $s \in \mathcal{D}$ such that $\|psp\| < 1 - \epsilon$.

I have been unable to falsify this version of the conjecture. I believe that a counterexample to this version would probably lead to a negative solution to the Kadison-Singer problem.

Since our counterexample is rather computational, it may be helpful to outline the geometric idea that underlies it. The plan is to produce a projection p and a unit vector v such that for any symmetry $s \in \mathcal{D}$ the quantity $\|psp(v)\|$ is larger than $2\delta_p$. We will take v to be in the range of p .

Now s is going to be a diagonal matrix whose entries are all ± 1 . So, letting $w_i = p(e_i)$ ($1 \leq i \leq n$), we will have

$$v = p(v) = p\left(\sum \langle v, e_i \rangle e_i\right) = \sum \langle v, w_i \rangle w_i$$

and

$$psp(v) = ps(v) = p\left(\sum \pm \langle v, e_i \rangle e_i\right) = \sum \pm \langle v, w_i \rangle w_i.$$

Thus, the vectors $\langle v, w_i \rangle w_i$ sum to the unit vector v , and the norm of $psp(v)$ for various symmetries s depends on the amount of cancellation which can be achieved by introducing sign changes in the sum.

The idea of the counterexample is to try to arrange p so that the vectors $w'_i = \langle v, w_i \rangle w_i$ approximate Figure 1. That is, all but one of them are identical, small, and nearly parallel to v , and the other one has a relatively large component perpendicular to v . Then regardless of how the symmetry s is chosen, the sum which constitutes $psp(v)$ can be bounded below in norm. For, if most of the small vectors are given the same sign as the large vector, then the sum will approximate $\pm v$, which has unit norm; but if most of the small vectors are given the opposite sign to v then the sum will have a large component perpendicular to v .

The actual counterexample is optimized by having not one vector with a large perpendicular component, but several, each perpendicular in a different direction. In the notation of the next section, these terms correspond to the basis vectors b_i . The small vectors nearly parallel to v correspond to the basis vectors a_i .

In our example the projection p is defined by specifying an orthonormal basis $\{v_k\}$ of its range, with v_0 playing the role of the vector v . The components of the v_k in the a_i and b_i directions are chosen so that the vectors $\langle v, w_i \rangle w_i$ will have approximately the form described above. In order to make the v_k orthonormal, and not increase δ_p , we also need to include further basis vectors c_{ij} and d_{ij} .

2. A counterexample to Conjecture A

Let $m \geq 6$ and let H be a complex Hilbert space of dimension $2m^3 + 8m^2 + 7m + 2$ (this number is not important). We fix an orthonormal basis for H whose members fall in four groups, labelled a through d . The basis vectors will be denoted a_i ($1 \leq i \leq m^2$), b_i ($1 \leq i \leq 2m+1$), c_{ij} ($1 \leq i < j \leq 2m+1$), d_{ij} ($1 \leq i \leq 2m+1$, $1 \leq j \leq (m+1)^2$). (Thus, the total dimension of H is

$$m^2 + (2m+1) + (2m^2 + m) + (m+1)^2(2m+1) = 2m^3 + 8m^2 + 7m + 2.)$$

$\mathcal{D}$ is the MASA of $B(H)$ consisting of those operators which are diagonal with respect to this basis.

We define p to be the orthogonal projection onto the $2m+2$ -dimensional subspace of H which is spanned by the following orthonormal set of vectors:

$$v_0 = \sum_{i=1}^{m^2} \frac{1}{m+1} a_i + \sum_{i=1}^{2m+1} \frac{1}{m+1} b_i$$

and for $1 \leq i \leq 2m+1$

$$\begin{aligned} v_i = & \sum_{j=1}^{m^2} \frac{-1}{m^2(m+1)} a_j + \frac{1}{m+1} b_i \\ & + \sum_{j=1}^{i-1} \frac{1}{m(m+1)} c_{ji} + \sum_{j=i+1}^{2m+1} \frac{-1}{m(m+1)} c_{ij} + \sum_{j=1}^{(m+1)^2} \frac{1}{m} \sqrt{\frac{m-1}{m+1}} d_{ij}. \end{aligned}$$

An easy computation shows that these vectors are indeed orthonormal.

Next, we show that $\delta_p = 2/(m+1)^2$. We have

$$\begin{aligned} \|p(a_i)\|^2 &= \left\| \frac{1}{m+1} v_0 + \sum_1^{2m+1} \frac{-1}{m^2(m+1)} v_i \right\|^2 = \frac{1}{(m+1)^2} + (2m+1) \cdot \frac{1}{m^4(m+1)^2} \\ \|p(b_i)\|^2 &= \left\| \frac{1}{m+1} v_0 + \frac{1}{m+1} v_i \right\|^2 = \frac{2}{(m+1)^2} \\ \|p(c_{ij})\|^2 &= \left\| \frac{-1}{m(m+1)} v_i + \frac{1}{m(m+1)} v_j \right\|^2 = \frac{2}{m^2(m+1)^2} \\ \|p(d_{ij})\|^2 &= \left\| \frac{1}{m} \sqrt{\frac{m-1}{m+1}} v_i \right\|^2 = \frac{m-1}{m^2(m+1)}. \end{aligned}$$

For $m \geq 6$ the largest of these is $2/(m+1)^2$, so this is the value of δ_p .

Now let s be any symmetry in $\mathcal{D}$, so that $s = q - q^\perp$ where q is a projection in $\mathcal{D}$. Let

$$\alpha = \#\{i : 1 \leq i \leq m^2, a_i \in \text{range}(q)\}.$$

By replacing s with $-s$ if necessary, we may suppose $\alpha \leq m^2/2$. For $1 \leq i \leq m^2$ say $s(a_i) = \epsilon_i a_i$ ($\epsilon_i = \pm 1$), and for $1 \leq i \leq 2m+1$ say $s(b_i) = \epsilon'_i b_i$ ($\epsilon'_i = \pm 1$).

We calculate $psp(v_0)$; its norm will give us a lower bound for $\|psp\|$:

$$\begin{aligned} psp(v_0) = & \left(\sum_1^{m^2} \frac{1}{(m+1)^2} \epsilon_i + \sum_1^{2m+1} \frac{1}{(m+1)^2} \epsilon'_i \right) v_0 \\ & + \left(\sum_1^{m^2} \frac{-1}{m^2(m+1)^2} \epsilon_i + \frac{1}{(m+1)^2} \epsilon'_1 \right) v_1 \\ & + \cdots + \left(\sum_1^{m^2} \frac{-1}{m^2(m+1)^2} \epsilon_i + \frac{1}{(m+1)^2} \epsilon'_{2m+1} \right) v_{2m+1}. \end{aligned}$$

Clearly the norm of $psp(v_0)$ exceeds the norms of its projections onto $\text{span}(v_0)$ and $\text{span}(v_1, \dots, v_{2m+1})$. So if $\alpha \leq m^2/4$ then (by projection onto $\text{span}(v_0)$)

$$\begin{aligned}
\|psp(v_0)\| &\geq \left| \sum_1^{m^2} \frac{1}{(m+1)^2} \epsilon_i + \sum_1^{2m+1} \frac{1}{(m+1)^2} \epsilon'_i \right| \\
&= \left| \frac{2\alpha - m^2}{(m+1)^2} + \sum_1^{2m+1} \frac{1}{(m+1)^2} \epsilon'_i \right| \\
&\geq \frac{m^2}{2} \cdot \frac{1}{(m+1)^2} - (2m+1) \cdot \frac{1}{(m+1)^2} \\
&\geq \frac{m^2 - 4m - 2}{4} \delta_p;
\end{aligned}$$

while if $\alpha \geq m^2/4$ then (by projection onto $\text{span}(v_1, \dots, v_{2m+1})$)

$$\begin{aligned}
\|psp(v_0)\| &\geq \sqrt{2m+1} \cdot \left| \sum_1^{m^2} \frac{-1}{m^2(m+1)^2} \epsilon_i \pm \frac{1}{(m+1)^2} \right| \\
&= \sqrt{2m+1} \cdot \left| \frac{m^2 - 2\alpha}{m^2(m+1)^2} \pm \frac{1}{(m+1)^2} \right| \\
&\geq \sqrt{2m+1} \cdot \frac{2\alpha}{m^2(m+1)^2} \\
&\geq \frac{\sqrt{2m+1}}{4} \cdot \delta_p.
\end{aligned}$$

We conclude that for any symmetry s in $\mathcal{D}$,

$$\begin{aligned}
\|psp\| &\geq \frac{\delta_p}{4} \cdot \min(m^2 - 4m - 2, \sqrt{2m+1}) \\
&= \delta_p \cdot \sqrt{m+1}/2
\end{aligned}$$

3. A limitation of the method

The counterexample presented in Section 2 is, in a sense, stronger than needed. To falsify the Akemann-Anderson conjecture, we had to place a lower bound on $\|psp\|$ for any symmetry $s \in \mathcal{D}$. That is, for any s , we had to find a unit vector v such that $\|psp(v)\|$ was sufficiently large. In fact, our counterexample was such that the same vector v_0 could play this role for all symmetries.

In pursuing stronger counterexamples, it is worth observing that there is a limitation to this approach. Namely, for any single vector v it is always possible to find a symmetry s

such that $\|psp(v)\|$ is on the order of $\delta_p^{1/2}$. This means that “single vector” counterexamples cannot falsify Conjecture B. In detail, we have the following result. We retain the notation of Section 1.

Theorem 1. *Let $p \in M_n(\mathbf{C})$ be a projection. For any unit vector $v \in \mathbf{C}^n$ there exists a symmetry $s \in \mathcal{D}$ such that*

$$\|psp(v)\| \leq \sqrt{2\delta_p + 3\delta_p^2}.$$

The proof of the theorem requires the following lemmas.

Lemma 2. *Let (v_i) be a sequence of vectors in a Hilbert space and let (w_i) be the sequence of partial sums, $w_i = v_1 + \cdots + v_i$. Suppose that*

$$\operatorname{Re}\langle v_{i+1}, w_i \rangle \leq 0$$

for all i . Then

$$\|w_i\|^2 \leq \|v_1\|^2 + \cdots + \|v_i\|^2$$

for all i .

Proof. By induction:

$$\begin{aligned} \|w_{i+1}\|^2 &= \|w_i\|^2 + \|v_{i+1}\|^2 + 2\operatorname{Re}\langle v_{i+1}, w_i \rangle \\ &\leq \|w_i\|^2 + \|v_{i+1}\|^2 \end{aligned}$$

and $\|w_1\|^2 = \|v_1\|^2$. ■

Lemma 3. *Let (v_i) be a finite sequence of vectors such that $\sum v_i = 0$. Then the sequence can be rearranged so as to satisfy the hypothesis of Lemma 2.*

Proof. Take v_1 for the first vector in the rearrangement. Then since $\sum_{i \neq 1} v_i = -v_1$, and hence

$$\left\langle \sum_{i \neq 1} v_i, v_1 \right\rangle \leq 0,$$

we must have

$$\operatorname{Re}\langle v_i, v_1 \rangle \leq 0$$

for some $i \neq 1$. Take this v_i for the second vector in the rearrangement, and continue inductively. ■

Proof of Theorem. It will suffice to consider the case that $v \in \operatorname{range}(p)$ (otherwise replace v by $p(v)$). Let $\{q_i\}$ be the minimal projections in $\mathcal{D}$, $q_i =$ projection onto $\operatorname{span}(e_i)$. Let p_1 be the projection onto $\operatorname{span}(v)$ and $p_2 = p - p_1$. For $1 \leq i \leq n$ let $\alpha_i = \|p_1(e_i)\|$,

$\beta_i = \|p_2(e_i)\|$, $x_i = p_1q_i p(v)$, and $y_i = p_2q_i p(v)$. Note that $x_i + y_i = pq_i p(v)$; $\|x_i\| = \alpha_i^2$ and $\|y_i\| = \alpha_i\beta_i$; and

$$\delta_p = \max\{\|p(e_i)\|^2\} = \max\{\alpha_i^2 + \beta_i^2\}.$$

Now $\sum pq_i p(v) = v$, so $\sum x_i = v$ and $\sum y_i = 0$. By Lemma 3 we may (rearranging the indices) suppose that the sequence (y_i) satisfies the hypothesis of Lemma 2. Since $\sum x_i = v$ and $x_i = \alpha_i^2 v$ it follows that $\sum \alpha_i^2 = 1$; also as $\alpha_i^2 \leq \delta_p$ for all i , we can find an integer k such that $|1/2 - \sum_1^k \alpha_i^2| \leq \delta_p/2$.

Let $q = q_1 + \cdots + q_k$ and let $s = q - q^\perp = 2q - 1$. Then by the choice of k we have

$$\|p_1 s p(v)\| = \|p_1 q(v) - p_1 q^\perp(v)\| = \left| \sum_1^k \alpha_i^2 - \sum_{k+1}^n \alpha_i^2 \right| \leq \delta_p,$$

and by Lemma 2,

$$\begin{aligned} \|p_2 s p(v)\|^2 &= \|p_2(2q - 1)p(v)\|^2 = 4\|p_2 q p(v)\|^2 = 4 \left\| \sum_1^k y_i \right\|^2 \\ &\leq 4 \sum_1^k \|y_i\|^2 = 4 \sum_1^k \alpha_i^2 \beta_i^2 \leq 4\delta_p \sum_1^k \alpha_i^2 \\ &\leq 4\delta_p(1/2 + \delta_p/2) = 2\delta_p + 2\delta_p^2. \end{aligned}$$

Thus

$$\|p s p(v)\|^2 \leq \delta_p^2 + 2\delta_p + 2\delta_p^2 = 2\delta_p + 3\delta_p^2,$$

as desired. ■

In light of this limitation, significant strengthening of the counterexample will require that the vector v be allowed to vary. The simplest way to achieve this would be to fix two vectors, v_0 and v_1 , at the outset, and arrange matters so that no matter how s is chosen either $\|p s p(v_0)\|$ or $\|p s p(v_1)\|$ is larger than some desired lower bound (the dichotomy depending, perhaps, on a parameter similar to α as in the example of Section 2). This would be sufficient to bound $\|p s p\|$ below, and it would allow greater flexibility in constructing examples. Using this method I have been able to construct a series of examples slightly improving those of Section 2, for which $\|p s p\|$ is bounded below by a constant times $\delta_p^{1/2}$.

[1] C. A. Akemann and J. Anderson, *Lyapunov Theorems for Operator Algebras*, *Mem. Amer. Math. Soc.* **94** no. 458 (1991).

[2] ———, The continuous Beck-Fiala theorem is optimal, *Discrete Math.* **146** (1995), 1-9.

- [3] J. Anderson, Extreme points in sets of positive linear maps on $B(H)$, *J. Funct. Anal.* **31** (1979), 195-217
- [4] J. Beck and T. Fiala, “Integer-making” theorems, *Discrete Appl. Math.* **3** (1981), 1-8.
- [5] J. Bourgain and L. Tzafriri, On a problem of Kadison and Singer, *J. Reine Angew. Math.* **420** (1991), 1-43.
- [6] R. V. Kadison and I. M. Singer, Extensions of pure states, *Amer. J. Math.* **81** (1959), 383-400.
- [7] G. A. Reid, On the Calkin representation, *Proc. London Math. Soc.* **23** (1971), 547-564.
- [8] B. Tanbay, Pure state extensions and compressibility of the l_1 -algebra, *Proc. Amer. Math. Soc.* **113** (1991), 707-713.

Math Dept.
 Washington University
 St. Louis, MO 63130
 nweaver@math.wustl.edu




Figure 1: Idea of the counterexample